

## PACKAGE LEAFLET: INFORMATION FOR THE USER

---

### ANGAL LEMON

5mg + 1mg, Compressed pastille

CHLORHEXIDINE, LIDOCAINE

---

*• This leaflet is a copy of the Summary of Product Characteristics and Patient Information Leaflet for a medicine, which outlines the conditions under which the medicine should be used and information on its known safety • The product information may be updated several times within its shelf life, and there could be differences between the version of information shown here and other information in the public domain or in the package insert • This leaflet may not contain all the information about the medicine or the information may not be the most up to date version for this product • If you have any questions or are not sure about anything, ask your doctor or pharmacist • Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.*

---

*• Keep this leaflet • You may need to read it again • If you have any further questions, ask your doctor or pharmacist • This medicine has been prescribed for you only • Do not pass it on to others • It may harm them, even if their signs of illness are the same as yours • If you get any side effects, talk to your doctor or pharmacist • This includes any possible side effects not listed in this leaflet •*

---

#### What is in this leaflet?

1. What ANGAL LEMON is and what it is used for
2. Before you take ANGAL LEMON
3. How to take ANGAL LEMON
4. Possible side effects
5. How to store ANGAL LEMON
6. Further information

#### 1. WHAT ANGAL LEMON IS AND WHAT IS IT USED FOR

ANGAL LEMON is a pastille with local effect in the mouth and pharynx.

ANGAL LEMON compressed pastille is antiseptic (local antimicrobial activity) and local painkiller.

ANGAL LEMON compressed pastille is recommended for:

- treatment of symptoms of sore throat, redness in the throat, discomfort in the throat accompanied with irritability
- in case of a bacterial infection accompanied with fever, additional treatment is necessary.

ANGAL LEMON compressed pastille contains no sugar and is therefore suitable for diabetics (people with diabetes).

#### 2. BEFORE YOU TAKE ANGAL LEMON

##### **Do not take ANGAL LEMON**

- If you are allergic to any ingredient of ANGAL LEMON pastille. All ingredients can be found in section 6.
- If your child is younger than 6 years.

**Take special care with ANGAL LEMON**

Talk to your doctor or pharmacist before using ANGAL LEMON pastille.

ANGAL LEMON pastille should not be used over a long period of time, without interruption or repeatedly. Limit its use to clearly necessary relief of pain and irritation.

In case of overdose (more than 20 pastilles per day) there is a small risk of impaired control over the swallowing reflex.

ANGAL LEMON pastille contains alcohol and propylene glycol. Please also read the section "ANGAL LEMON compressed pastille contains alcohol and propylene glycol."

If you use other medicines, please read the section "Taking other medicines".

Contact your doctor if any of the above applies to you or if anything like that applied to you in the past.

**Children**

ANGAL LEMON pastille should not be used in children younger than 6 years.

**Taking other medicines**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines obtained without a prescription and herbal medicines. Do not use ANGAL LEMON pastille with medicines for the treatment of mild forms of Alzheimer's disease (gradual and permanent loss of memory, orientation and language capability), such as cholinesterase inhibitors (medicines for treating dementia).

During the administration of ANGAL LEMON you should not simultaneously use other antiseptics.

**Pregnancy and breast-feeding**

Before you start using ANGAL LEMON pastille, talk to your doctor or pharmacist if you are pregnant, planning to become pregnant or are breastfeeding.

ANGAL LEMON pastille contains alcohol. Please read part "ANGAL LEMON compressed pastille contains alcohol, and propylene glycol."

If you follow previously mentioned precautions, ANGAL LEMON compressed pastille may be used during pregnancy or breastfeeding. Tell your doctor that you are using this medicine.

**Driving and using machines**

It is not likely that ANGAL LEMON will affect your ability to drive and use machines.

**Other warnings**

ANGAL LEMON compressed pastilles contain sorbitol. If your doctor ever said that you have intolerance to certain sugars, consult your doctor before using this medicine. Sorbitol can have a mild laxative (stool softener) effect. The caloric value is 2.6 kcal / g of sorbitol.

ANGAL LEMON compressed pastilles also contain aspartame. It is a source of phenylalanine and may be harmful for people with phenylketonuria (a metabolic disorder that is characterized by elevated levels of phenylalanine / amino acid / in the blood).

**3. HOW TO TAKE ANGAL LEMON**

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure.

**Recommended dosage:**

Adults: 6 to 10 compressed pastilles a day.

Allow the compressed pastille to slowly dissolve in the mouth.

Children older than 6 years: 3-5 compressed pastilles a day.

Allow the compressed pastille to slowly dissolve in the mouth.

This medicine should not be used over long periods of time. Contact your doctor if symptoms worsen or do not improve after 3-4 days.

***If you use more ANGAL LEMON than you should***

If you take more of this medicine than you should, talk to a doctor or go to a hospital straight away. This medicine contains only small portion of the toxic dose. If used properly, the risk of overdose is very small. Symptoms of overdose are: restlessness, tinnitus, breathing difficulties, cardiac arrhythmia (abnormal rhythm of the heart) and reduced control over the swallowing reflex.

Talk to your doctor or pharmacist if you have any additional questions.

#### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, **ANGAL LEMON** can cause side effects, although not everybody gets them. Side effects are classified according to frequency: very common: 1 / 10; common: (1/10 - 1/100), uncommon (1/100 - 1/1000); rare (1 / 1,000 to 1 / 10,000); very rare (1 / 10,000), Not known: frequency cannot be determined from the available data.

When used in the recommended manner, ANGAL LEMON pastille is well tolerated.

**Very rare:** allergic reactions on the skin and mucous membranes.

In special cases it may cause changes to the senses of taste and anaphylactic reactions (hypersensitivity reactions).

In the case of prolonged and continuous application you may experience brown coloration of teeth. However, this brown color is easy to remove.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **5. HOW TO STORE ANGAL LEMON**

Keep out of the reach and sight of children.

Do not store above 25 °C.

Do not use **ANGAL LEMON** after the expiry date which is stated on the label, 1 year from the date of manufacture.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### **6. FURTHER INFORMATION**

***What ANGAL LEMON contains***

The active substances are chlorhexidine hydrochloride 5 mg and lidocaine hydrochloride 1 mg.

The other ingredients are:

sorbitol

magnesium stearate

aspartame

lemon flavor 5051050 AP0551

Acesulfame-K

***What ANGAL LEMON looks like and contents of the pack***

ANGAL LEMON compressed pastilles are available in a box with 24 compressed pastilles.

***Regime of dispensing***

The medicine is issued without doctor's prescription.

***Manufacturer***

Lek farmacevtska družba d.d., Verovškova 57, Ljubljana, Slovenia

***Manufacturer of the medicinal product***

Laboratoria Qualiphar N.V./S.A., Rijksweg 9, B-2880 Bornem, Belgium

Lek farmacevtska družba dd, Veroškova 57, Ljubljana, Slovenia