

PACKAGE LEAFLET: INFORMATION FOR THE USER

ATRICAN

250 mg gastro-resistant capsule, soft

TENONITROZOLE

• This leaflet is a copy of the Summary of Product Characteristics and Patient Information Leaflet for a medicine, which outlines the conditions under which the medicine should be used and information on its known safety • The product information may be updated several times within its shelf life, and there could be differences between the version of information shown here and other information in the public domain or in the package insert • This leaflet may not contain all the information about the medicine or the information may not be the most up to date version for this product • If you have any questions or are not sure about anything, ask your doctor or pharmacist • Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

• Keep this leaflet • You may need to read it again • If you have any further questions, ask your doctor or pharmacist • This medicine has been prescribed for you only • Do not pass it on to others • It may harm them, even if their signs of illness are the same as yours • If you get any side effects, talk to your doctor or pharmacist • This includes any possible side effects not listed in this leaflet •

What is in this leaflet?

1. What ATRICAN tablets is and what it is used for
2. Before you take ATRICAN
3. How to take use ATRICAN
4. Possible side effects
5. How to store ATRICAN
6. Further information

1. WHAT ATRICAN IS AND WHAT IT IS USED FOR

This medicine is prescribed for the treatment of urogenital trichomoniasis, which is an infection caused by the parasite *Trichomonas vaginalis* (the infection of the genital tract of women and men).

2. BEFORE YOU TAKE ATRICAN

Do not take ATRICAN

- If you are allergic (hypersensitive) to Tenonitroazole or to any of the other ingredients in this medicine
- If you have liver failure
- If you are allergic to peanuts or soya

Take special care with ATRICAN, so you must tell the doctor if you

Warning:

This medicine should not be taken with alcohol.

Precautions:

- Tell your doctor, in case of hematological disorders associated with white blood cells,
Do not wear contact lenses during treatment.

- Your Doctor may also treat you and your partner.

If your doctor has told you that you have an intolerance to some sugars, before you start taking this medicine, consult your doctor.

Taking ATRICAN with food and drink

ATRICAN should not be taken with alcohol.

Pregnancy and breastfeeding

The medicinal product ATRICAN contains a substance (dibutyl phthalate) which can cause disorders of the endocrine system, in the amount exceeding the permitted limits.

For this reason, the use of ATRICAN is not recommended during pregnancy and breastfeeding.

Important information about some of the ingredients in ATRICAN

List of other substances that are known to have confirmed effect: sorbitol, soybean oil, peanut oil.

3. HOW TO TAKE ATRICAN**For adults only.**

1 capsule in the morning and in the evening during meals, 4 days.

Oral use.

Both partners should be treated simultaneously.

If you take more ATRICAN than you should

If you take more ATRICAN than you should immediately talk to your doctor or pharmacist.

If you forget to take ATRICAN

Do not take a double dose to make up for a forgotten dose.

Talk to your doctor or pharmacist if you have any additional questions.

4. POSSIBLE SIDE EFFECTS

Like all medicines, ATRICAN can cause side effects, although not everybody gets them.

- mild stomach problems: nausea, feeling of heaviness in the stomach, loss of appetite
- Staining of contact lenses in yellow

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting suspected side effects

In case of occurrence of any side effects after administration of the medicine, it is necessary to inform your doctor or pharmacist. This includes all possible side effects that are not included in this leaflet, as well as those that are.

5. HOW TO STORE ATRICAN

Keep out of the reach and sight of children.

Do not store above 25 °C.

Do not use ATRICAN after the expiry date which is stated on the label, 3 years from the date of manufacture.

6. FURTHER INFORMATION

What ATRICAN contains

1 gastro-resistant capsules, soft, contains 250 mg of the active substance tenonitroazole.

Other ingredients: peanut oil, hydrogenated soybean oil, soya lecithin, gelatin, sorbitol and sorbitan solution, titanium dioxide (E 171), iron oxide (E 172), yellow, glycerol, hypromellose phthalate, dibutyl phthalate.

What ATRICAN looks like and contents of the pack

Oval, yellow to yellow-brown soft capsule, containing a homogeneous oily mass.

The outer packaging of the drug: a cardboard box with a blister which contains 8 capsules.

Regime of dispensing

The medicine is issued on prescription.

Manufacturer

Laboratoire Innotech International
22 avenue Aristide Briand
Arcueil, France

Manufacturer of the medicinal product

Innothera Chouzy
Rue René Chantereau-L'Isle Vert
Chouzy-sur-Cisse
France